Table 1. Number, incidence rate <sup>1</sup>, median days away from work <sup>2</sup> and relative standard errors <sup>3</sup> of occupational injuries and illnesses involving days away from work <sup>4</sup> by selected natures with musculoskeletal disorders <sup>5</sup> in private industry for Nebraska, 2005

Nature of the injury or illness	Total Cases	Incidence Rate	Median Days	Relative Standard Error
All Natures	3,090	50.5	6	4.9
021 Sprains- strains- tears	2,030	33.3	5	5.4
0972 Back pain- hurt back	230	3.8	5	11.8
0973 Soreness- pain- hurt- except the back	190	3.2	19	12.9
1241 Carpal tunnel syndrome	290	4.7	8	10.8
153 Hernia	180	2.9	18	13.3
1530 Hernia- unspecified	30	0.6	18	29.5
1531 Inguinal hernia	90	1.5	19	18.2
1533 Ventral hernia	50	0.8	18	25.2
17 Musculoskeletal system and connective tissue diseases and disorders	160	2.6	6	14.2
173 Rheumatism- except the back	150	2.4	7	14.6
1731 Bursitis	20	0.3	12	37.3
1733 Tendonitis	20	0.4	34	35.0
1739 Rheumatism- except the back- n.e.c.	80	1.3	3	19.4

<sup>1</sup> Incidence rates represent the number of injuries and illnesses per 10,000 full-time workers and were calculated as: (N / EH) X 20,000,000 where,

N = number of injuries and illnesses,

EH = total hours worked by all employees during the calendar year,

20,000,000 = base for 10,000 full-time equivalent workers (working 40 hours per week, 50 weeks per year).

NOTE: Dashes indicate data that do not meet publication guidelines or data for incidence rates less than .05 per 10,000 full-time workers. The scientifically selected probability sample used was one of many possible samples, each of which could have produced different estimates. A measure of sampling variability for each estimate is available upon request.

SOURCE: Bureau of Labor Statistics, U.S. Department of Labor, November 2006

<sup>&</sup>lt;sup>2</sup> Median days away from work is the measure used to summarize the varying lengths of absences from work among the cases with days away from work. Half the cases involved more days and half involved less days than a specified median. Median days away from work are represented in actual values.

<sup>&</sup>lt;sup>3</sup> Relative standard errors are a measure of the sampling error of an estimate. Sampling errors occur because observations are made on a sample, not on the entire population. Estimates based on the different possible samples of the same size and sample design could differ. Relative standard errors less than 0.05 are not shown.

<sup>&</sup>lt;sup>4</sup> Days-away-from-work cases include those that result in days away from work with or without job transfer or restriction.

<sup>&</sup>lt;sup>5</sup> Includes cases where the nature of injury is: sprains, strains, tears; back pain, hurt back; soreness, pain, hurt, except back; carpal tunnel syndrome; hernia; or musculoskeletal system and connective tissue diseases and disorders and when the event or exposure leading to the injury or illness is: bodily reaction/bending, climbing, crawling, reaching, twisting; overexertion; or repetition. Cases of Raynaud's phenomenon, tarsal tunnel syndrome, and herniated spinal discs are not included. Although these cases may be considered MSD's, the survey classifies these cases in categories that also include non-MSD cases.